arXiv:0707.4076v1 [nucl-th] 27 Jul 2007

Thermal model predictions of hadron ratios at LHC

A Andronic[1], **P Braun-Munzinger**[1,2], **J Stachel**[3]

[1] Gesellschaft für Schwerionenforschung, GSI, D-64291 Darmstadt, Germany
[2] Technical University Darmstadt, D-64283 Darmstadt, Germany
[3] Physikalisches Institut der Universität Heidelberg, D-69120 Heidelberg, Germany

Abstract. We present predictions of the thermal model for hadron ratios in central Pb+Pb collisions at LHC.

Based on the latest analysis within the thermal model of the hadron yields in central nucleus-nucleus collisions [1], the expected values at LHC for the chemical freeze-out temperature and baryochemical potential are T=161±4 MeV and μ_b=$0.8^{+1.2}_{-0.6}$ MeV, respectively. For these values, the thermal model predictions for hadron yield ratios in central Pb+Pb collisions at LHC are shown in Table 1. We have assumed no contribution of weak decays to the yield of pions, kaons and protons.

Table 1. Predictions of the thermal model for hadron ratios in central Pb+Pb collisions at LHC. The numbers in parantheses represent the error in the last digit(s) of the calculated ratios.

π^-/π^+	K^-/K^+	$\bar{p}/p$	$\bar{\Lambda}/\Lambda$	$\bar{\Xi}/\Xi$	$\bar{\Omega}/\Omega$
1.001(0)	0.993(4)	$0.948^{-0.013}_{+0.008}$	$0.997^{-0.011}_{+0.004}$	$1.005^{-0.007}_{+0.001}$	1.013(4)

p/π^+	K^+/π^+	K^-/π^-	Λ/π^-	Ξ^-/π^-	Ω^-/π^-
0.074(6)	0.180(0)	0.179(1)	0.040(4)	0.0058(6)	0.00101(15)

The antiparticle/particle ratios are all very close to unity, with the exception of the ratio $\bar{p}/p$, reflecting the expected small, but nonzero, μ_b value. The errors are determined by the errors of μ_b in case of antiparticle/particle ratios and by the errors of T for all other ratios.

Table 2. Predictions for the relative abundance of resonances at chemical freeze-out.

ϕ/K^-	K^{*0}/K^0_S	Δ^{++}/p	$\Sigma(1385)^+/\Lambda$	Λ^*/Λ	$\Xi(1530)^0/\Xi^-$
0.137(5)	0.318(9)	0.216(2)	0.140(2)	0.075(3)	0.396(7)

Assuming that the yield of resonances is fixed at chemical freeze-out, we show in Table 2 predictions for the relative yield of various resonance species. We emphasize that the above hypothesis needs to be checked at LHC, in view of the data at RHIC [2], which may indicate rescattering and regeneration of resonances after chemical freeze-out.

References

[1] A. Andronic, P. Braun-Munzinger, and J. Stachel. Hadron production in central nucleus nucleus collisions at chemical freeze-out. *Nucl. Phys.*, A772:167–199, 2006.

[2] John Adams et al. Strange baryon resonance production in s(nn)**(1/2) = 200- gev p + p and au + au collisions. *Phys. Rev. Lett.*, 97:132301, 2006.